



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

The diversity of maternal-age effects upon pre-adult survival across animal species

Citation for published version:

Ivimey-Cook, E & Moorad, J 2020, 'The diversity of maternal-age effects upon pre-adult survival across animal species', *Proceedings of the Royal Society B: Biological Sciences*, vol. 287, no. 1932, 20200972. <https://doi.org/10.1098/rspb.2020.0972>

Digital Object Identifier (DOI):

[10.1098/rspb.2020.0972](https://doi.org/10.1098/rspb.2020.0972)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Proceedings of the Royal Society B: Biological Sciences

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



The diversity of maternal-age effects upon pre-adult survival across animal species

Edward Ivimey-Cook^{1*} and Jacob Moorad¹

¹Institute of Evolutionary Biology, School of Biological Sciences, University of Edinburgh, Edinburgh, EH9 3JT, UK

Abbreviated Title

Diversity of maternal-age effects

Corresponding author

E.Ivimey-Cook@uea.ac.uk

Keywords

Ageing, maternal-effect senescence, meta-analysis, demography, pre-adult, survival

Abstract

Maternal senescence is the detrimental effect of increased maternal age on offspring performance. Despite much recent interest given to describing this phenomenon, its distribution across animal species is poorly understood. A review of the published literature finds that maternal age affects pre-adult survival in 252 of 272 populations (93%) representing 97 animal species. Age effects tended to be deleterious in invertebrates and mammals, including humans, confirming the presence of senescence. However, bird species were a conspicuous exception, as pre-adult survival tended to increase with maternal age in surveyed populations. In all groups, maternal-age effects became more negative in older mothers. Invertebrates senesced faster than vertebrates, and humans aged faster than non-human mammals. Within invertebrates, Lepidopterans demonstrated the most extreme rates of maternal-effect senescence. Among the surveyed studies, phylogeny, life history, and environment (e.g., laboratory vs wild populations) were tightly associated; this made it difficult to make confident inferences regarding the causes of diversity for the phenomenon. However, we provide some testable suggestions, and we observe that some differences appear to be consistent with predictions from evolutionary theory. We discuss how future work may help clarify ultimate and proximate causes for this diversity.

Introduction

Senescence is the age-related physiological deterioration of organismal function typically associated with increasing mortality risk (actuarial senescence) and decreasing fertility (reproductive senescence). Studies report actuarial and reproductive senescence in most animal species across most phyla [1–6], with especially well documented senescent declines in wild vertebrates [7–10] and laboratory invertebrates [11–14]. Maternal senescence, the detrimental result of a mother's increasing age on traits associated with offsprings' life history or fitness [15–19], is a distinctly different manifestation of age. Whilst such effects of maternal age are attracting increased scientific attention, their distributions across the tree-of-life remain poorly described [20]. Investigating the prevalence and degree of maternal-age effects is an important first step to understanding ultimate and proximate causes of this form of senescence, as this may identify taxa that have unusual manifestations of ageing that warrant special focus in the future. To clarify, we use the term 'ageing' to refer to any age-related change, but we reserve 'senescence' to indicate a deleterious effect of increased age.

Several hundreds of models have been proposed to explain the proximate causes of senescence [21–25]. In contrast, there are few explanatory evolutionary models, but all share the central tenet that senescence is caused ultimately by age-related declines in the efficacy of natural selection [26]. Mutation accumulation [27] and antagonistic pleiotropy [28] are two such models that make different assumptions regarding the genetic architecture of age-specific traits. Population genetic models use estimates of vital rates (age-specific survival and reproduction rates) and various assumptions related to gene action to predict patterns of actuarial senescence (e.g. [29,30]). More recently, Moorad and Nussey [31] modified these to quantify how age changes the strength of selection for age-specific maternal effects and to show how these changes cause maternal senescence

manifested upon pre-adult survival to evolve. They predicted that evolved demographic patterns of this senescence should be qualitatively different from actuarial and reproductive senescence. However, we know little about how well this model predicts patterns of ageing in real populations.

In this paper, we address conspicuous gaps in our understanding of the taxonomic breadth and intensity of maternal-effect ageing by performing an extensive systematic review of the literature using meta-analytical methodology. We chose pre-adult survival, defined here as survival throughout some part of the pre-reproductive period. The nature of this part will vary according to the methodologies of the available papers, and it largely reflects the characteristics of the study species (e.g., hatching rate in invertebrates, survival to fledging in birds, survival to weaning in mammals or child survival in humans). This trait was chosen as our focus for maternal-age effects for several reasons: 1) this trait's relationship to fitness is profound and well-understood conceptually [26]; 2) evolutionary theory explicitly models age-specific maternal effects on this trait [31]; and 3) associations between the trait and maternal age are observed with sufficient frequency to enable a large-scale review. This study addresses questions about the nature of maternal-effect ageing as it manifests on pre-adult survival rates:

1. *Does maternal age tend to affect pre-adult survival in most species?*
2. *Do effects of increased maternal age tend to be negative (is maternal senescence the norm)?*
3. *What features of specific studies (for example, phylogeny, presence/absence of biparental care, nature of human interventions) appear to predict effect sizes?*

We find that maternal-age effects are widespread across studies of animal species. However, senescence appears to be a general and important phenomenon in only some groups, with large observable variation in the rates of senescence across groups. Wild

birds and Lepidopterans (butterflies and moths) represent two disparate extremes. Why these particular taxonomic groups should be unusual in this respect is an ecological and evolutionary puzzle.

Materials and Methods

This meta-analysis followed the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (“PRISMA”) guidelines [32] (Fig. S1). A literature search was conducted in December 2019 using the online databases Web of Science and Scopus. Search terms are provided in Supplementary Table S1.

Accepted papers include the number of surviving and dying pre-adults as functions of maternal age (Fig. S1). Where a study was replicated within a paper or where a species was studied in more than one paper, discrete binomial datasets were extracted for each replicate population, and we treated all such within-species replicates as independent in subsequent analyses. Our criteria for acceptance and methods for data extraction are described in the Supplemental Methods. Each age class was associated with a corresponding number of surviving and dying pre-adults (coded with 1s and 0s, respectively) reconstructed from the realised maternal age distribution, the mean rates of age-specific fecundity, and pre-adult survival rates extracted from the source papers.

Maternal ages in each study were standardized by dividing by generation time T . For each replicate i , generation time T_i was calculated as the average of the maternal age distribution $f(x)$, or $T_i = \sum_x xN_{xi}/\sum_x N_{xi}$ [33]. However, this measure is sensitive to the age structure and vital rates of the population. In populations where the timing of breeding is influenced by experimenters who may have wished to enhance the power of a study to detect age-related effects, the value of T in the experimental population may not reflect the distribution of maternal ages found in natural or equilibrium populations.

Such cases likely involve the over-sampling of older ages; this would tend to inflate values of T and therefore underestimate the true magnitudes of maternal effects. Laboratory studies are most likely to introduce such biases. Failure to properly incorporate the duration of the pre-reproductive period into generation time calculations, which is often substantial in laboratory invertebrates, can also contribute to significant bias. Therefore, for studies where age was defined in terms of elapsed time since reaching adulthood, total age x was taken as the sum of adult age and the duration of the juvenile period. The latter was taken either from the source paper or from secondary sources.

All else equal, we expect that the rates of survival over some time interval will decrease as the size of that interval increases. Wherever possible, we extracted estimates of this pre-adult study interval, t , from each study. This study interval is standardized by generation length. For example, survival to one year of age was assessed in the olive baboon (*Papio anubis*), which has a generation time of 11.16 years (see [34] for estimates of T and pre-adult study duration). Here, $t = 1 \text{ yr} / 11.16 \text{ yr} = 0.09$. Some studies did not provide clear descriptions of study intervals. For example, time-to-hatch in *Drosophila melanogaster* eggs was unspecified in [35]. In cases such as these, an approximate study interval for that species was obtained from available secondary sources (e.g., 13.50 hours for *D. melanogaster* [36]).

Each study was assigned to one of four environmental groups: wild, laboratory, semi-captive (where humans provided food or veterinary intervention), and agricultural. Humans were also surveyed but not so categorized. Non-human animals were grouped by taxonomic relationships into invertebrates, birds, non-human mammals, and other vertebrates. Invertebrates were further divided into Lepidopterans and other invertebrates (including non-Lepidopteran insects), but this division was made after effect sizes were estimated, and a pronounced difference was observed between these

two groups. Phylogenetic trees were created using the National Centre for Biotechnology Information Taxonomy database [37] and PhyloT [38] and visualised using 'ggplot2' and 'ggtree' [39,40].

We estimated the effect that maternal age had on the proportion of surviving pre-adults for each replicate independently. We fit generalised linear models (GLMs) of pre-adult survival (P) with binomial error (e) distribution and probit link functions to: [1] age-independent, [2] linear and [3] quadratic models of maternal age. This link function assumes a Gaussian distribution of a latent predictor variable, and it is a standard function used by quantitative geneticists for scaling genetic contributions to survival [41,42]. For equations 1-3, $P(x)$ is the probability of a pre-adult surviving at standardised maternal age x , whereas A , B , and C are the intercept, the linear coefficient, and the quadratic coefficient, respectively.

$$P(x) = A + e \quad [1]$$

$$P(x) = A + Bx + e \quad [2]$$

$$P(x) = A + Bx + Cx^2 + e \quad [3]$$

Replicate-specific log-likelihoods for all models were noted along with estimates of effect sizes and associated standard errors (SEs). We calculated Akaike Information Criterion values (AIC) for each replicate i , and model j using $AIC_{ij} = 2k_j - 2\loglik_i$, where k_j is the number of parameters (one, two or three, for the age-independent, linear or quadratic models, respectively). From these, sample-size corrected AIC values ($AICc$) were calculated using the formula $AICc_{ij} = \frac{AIC_{ij} + 2k_j(k_j + 1)}{(n_i - k_j - 1)}$, where n_i was the number of observations for each replicate [43]. Based upon the results from model-fitting (see

Results), our subsequent analyses focus upon the linear effects of maternal age in old mothers, where “old” is defined as equal or greater than the average age of mothers T (standardized maternal ages greater than one).

To correct for variation in study intervals t across studies, we first assessed its impact on maternal-age effects. We performed a bootstrap weighted regression using the “boot” package in R Version 3.6.0 [44–46] of estimated linear effect sizes upon t . Weights were taken from the inverse of the estimated SEs that were associated with the linear effect sizes. Values of t are mathematically constrained on the interval $1 \geq t > 0$, and they ranged here from 0.007 in a human population to 0.935 in a Dermestid beetle species, *Trogoderma inclusum* (See Fig. S2 for a full distribution of t across all extracted populations). In general, invertebrates appeared to spend a greater fraction of their generation time observed as pre-adults than the vertebrates (median $t = 0.120$ and 0.091 , respectively). Note that two replicates of rotifers (*Brachionus calcyflorus*) included a diapause phase as part of the pre-adult period [47]. As this period of diapause described in the paper appeared to us to be both arbitrary and highly influential to calculated values of t , we considered only the time from egg to hatching and omitted the diapause phase from analysis of these populations.

The overall effect of t (pre-adult study interval) on maternal-age effects tended to be negative, -1.235 ($-2.750, 0.107$) [results presented in this way express the bootstrapped mean and the bias-corrected 95%-tiles in brackets]. Because greater values of t indicate greater exposure to mortality risk, and some risk can follow from maternal age, we expect that any average effect that age tends to have on survival across all studies will be amplified as t increase. As these age effects tended to be negative (see Result below), the negative influence of t upon those estimates is to be expected. We corrected age effects for the estimated effect of study interval by adding $1.235 \times t$ to each. We did not alter SEs

associated with these estimates. Replicates were pooled in groups in accordance to taxonomy and environments. From these groups, we calculated weighted bootstrapped means for each ($n = 10,000$ replicates), where weightings were the inverse of the estimated SEs.

Results

A total of 196 animal papers met our search criteria. From these, we extracted and analysed 273 populations from 97 animal species (Table S2). One population was disregarded because no offspring died within the experimental replicate. Species were studied in a single environmental context, with three exceptions: 1) the red-legged partridge, *Alectoris rufa*, (semi-captive and agricultural [48,49]); 2) the domestic sheep, *Ovis aries*, (agricultural and natural [50,51]); and 3) the Columbian ground squirrel *Spermophilus columbianus* (semi-captive and natural [52,53]).

There were strong associations among the available studies between taxonomic and environmental groups (Table 1). Invertebrate species were only studied in the laboratory. Birds and mammals were most frequently studied in the wild, and only one study from one species from either of these groups (*Mus musculus*) provided suitable laboratory data. Ideally, taxonomic groups would have been distributed more evenly over environments, as this might have supported a two-factor analysis. Unfortunately, these strong associations among the available literature undermine rigorous attempts to assign causes of potential effect size differences to either phylogeny (at the vertebrate/invertebrate scale) or environmental context. Compounding this problem is the fact that life history strategies share this same dichotomous partition of species: birds and mammals are long-lived and tend to provide obvious post-natal maternal care. In

contrast, the invertebrate species studied here in the laboratory are short-lived and demonstrate little or no conspicuous maternal care. The sole mammalian species to be studied in the laboratory is the only species studied in that context to provide post-natal maternal care. The only studied reptile species, *Lacerta vivipara*, provides pre-natal maternal care.

Replicate-specific results from the GLMs (Equations 1-3) are given in Table S3a. Comparisons of AICc values found that the age-independent models were best in 20 cases, linear age effect models were best in 51 cases, and quadratic age effect models were best in 201 cases. Summed AICc values over all replicates indicated a strong preference for the quadratic model of maternal age on pre-adult survival (ΔAICc Age-Independent: +229755; ΔAICc Linear: +43828). We estimated negative quadratic effects in 202 cases and positive quadratic effects in 70 cases. The weighted bootstrapped means of the quadratic effects were negative when pooled over all species -0.525 (-0.813, -0.308) and within all taxonomic groups (Table 1). Quadratic effects were different from zero in the aggregate and in all but two groups (birds and mammals). Nevertheless, the strong tendency towards a negative quadratic effect of age across species indicates that linear fits of all available maternal ages tend to underestimate senescence experienced by older females (or overestimate maternal effect improvement in the old). In light of this finding, we re-focused our question to evaluate the linear effects of maternal age on old females only, where old defines all ages greater than T (generation time). The distribution of age-effects in old mothers is illustrated in Fig. 1, and the range of ages that remained for each study after removing ages less than T are illustrated in Fig. S3. Note that the approach taken here to focus upon particular ages does not presume that senescence actually exists in any population; in this way it differs from other approaches that

estimate an age-of-onset for senescence and use this age to define ageing rates in identified senescence phases of life [54–56].

When averaged over all populations, the linear effects of maternal ages across all available ages were negative, -0.378 (-0.573, -0.204). The linear effects of ages *in the old*, which are the focus of our analyses, were stronger and remained statistically significant, -0.691 (-0.913, -0.505). All effect sizes are reported on the probit scale. Conversion to the scale of survival is not straightforward, as linearity on the probit scale implies strong nonlinearity on the scale of survival. Nonetheless, we provide one metric that can indicate the effects of increased maternal age at the onset of old age on this scale. We define $\delta(0.01 \times k)$ to be the increase in age from T that delivers a 1% change in pre-adult survival, where k is +1 (a 1% increase in survival) when age effects are positive ($B > 0$) and $k = -1$ (a 1% decrease in survival) with maternal senescence ($B < 0$). Smaller values indicate less time required to make that change and stronger age effects. These values are estimated for all replicates and reported in Table S3b and Fig. S4 of the supplementary material. The reported taxonomic structure is based upon probit measures, but the diversity illustrated in Fig S4 suggests that these qualitative patterns are robust to this change in scaling.

Populations were pooled within each environment-by-taxon group, and the bootstrapped means and 95%-tiles are reported for each in Table 1. Of the five most populated groups, all appeared to have mean effect sizes that differed from zero. Lepidopterans, other invertebrates, wild mammals, and humans exhibited statistically significant deleterious effects of maternal age. In contrast, wild birds appeared to present positive age effects on early survival. Senescence was most pronounced within the Lepidopterans, with deleterious age effects in the old of nearly an order of magnitude greater than the global mean (-6.142 vs -0.691).

269

270 **Table 1.** Maternal-age effects in the old for all environment-by-taxon groups (means and
 271 bias-corrected 95%-tiles). Sample sizes are given in italics where the number of species
 272 is followed by the number of replicates. Confidence intervals are not indicated when only
 273 one replicate was available. Bold-faced estimates of the means indicate significance at
 274 $\alpha < 0.05$.

	<i>Lepidopterans</i>	<i>Other Invertebrates</i>	<i>Birds</i>	<i>Mammals</i>	<i>Other Vertebrates</i>	<i>Humans</i>
<i>Laboratory</i>	-6.142 (-8.885, -4.088) <i>15/27</i>	-0.849 (-1.295, -0.471) <i>34/79</i>	- 	3.280 <i>1/1</i>	0.075 <i>1/1</i>	-
<i>Semi-Captive</i>		-	-0.515 (-0.937, 0.029) <i>4/4</i>	-0.228 (-0.723, 0.045) <i>7/8</i>	-0.986 <i>1/1</i>	-
<i>Agricultural</i>		-	0.327 (0.071, 1.246) <i>2/8</i>	0.137 (-0.258, 0.583) <i>3/13</i>	-	-
<i>Wild</i>		-	0.124 (0.002, 0.287) <i>20/37</i>	-0.295 (-0.451, -0.140) <i>11/12</i>	-	-
<i>Humans</i>		-	-	-	-	-0.819 (-1.113, -0.423) <i>1/80</i>

275

276 Finally, we paired four groups and assemblages of groups to compare effect sizes using
 277 Mann-Whitney U Tests. Two such assemblages are ‘Non-Human Mammals’ ($n = 34$) and
 278 All Vertebrates’ ($n = 165$). The bootstrapped means values for these are 0.0001 (-0.218,
 279 0.359) and -0.303 (-0.456, -0.164), respectively. Non-directional effects within the ‘Non-
 280 Human Mammals’ group is caused by combining positive effects from agricultural studies
 281 with negative effects from other mammal populations. The following comparisons of
 282 mean effect sizes follow from Table 1 and results from four independent Mann-Whitney
 283 Tests:

- 284 1. Lepidopterans < Non-Lepidopteran Invertebrates < 0 ($W = 198$, $P < 0.001$);
- 285 2. Humans < Non-Human Mammals ($W = 532$, $P < 0.001$);

3. Wild Mammals $< 0 <$ Wild Birds ($W = 64$, $P < 0.001$); and

4. Non-Lepidopteran Invertebrates $<$ All Vertebrates < 0 ($W = 5501$, $P = 0.049$).

Discussion

Our results provide definitive answers to two study goals. First, maternal age affected pre-adult survival rates in 93% of extracted populations drawn from divergent animal taxa and environments. Second, these effects tended to be deleterious (thereby fitting the definition of maternal-effect senescence) across all broadly defined animal groups, with the conspicuous exception of birds from agricultural and wild populations. Maternal age trajectories tended towards concavity in all groups, indicating that rates of senescence intensified in old mothers in populations that senesce, and rates of improvement diminished in late life in those populations that do not senesce.

The general trends observed here are anticipated by recent evolutionary theory. Moorad and Nussey [31] integrated *Indirect Genetic Effects* (IGEs) into evolutionary demographic models of phenotypic selection with the aim to predict how maternal age should evolve to affect pre-adult survival. IGEs are a quantitative genetic concept developed by animal breeders [57,58] before gaining much attention from evolutionary geneticists interested in social evolution [59–61]. This begins with the conventional perspective that individual phenotypes (e.g., pre-adult survival) are affected by their own genes (*Direct Genetic Effects*, or DGEs) and the environment that they experience. However, it also recognises that one's environment can be affected by influences from social partners (e.g., mothers), and the social environment that produces the phenotype can evolve by natural selection to the degree that these influences are genetic (IGEs).

309 Phenotypes evolve as individuals' genetically-determined environments change by
310 natural selection.

311 Most evolutionary genetic models of senescence assume implicitly that DGEs
312 represent the only route towards evolutionary change [26,28,62], but Moorad and
313 Nussey [31] modified one such model [30] by assuming that mothers contributed IGEs
314 that were independent and identically distributed across all ages. They found that as
315 maternal age increases, selection to remove deleterious age-specific IGEs must eventually
316 diminish, and it follows that the pre-adult survival should evolve such that it deteriorates
317 as mothers get older. Furthermore, they showed that certain demographic conditions
318 exist that allow pre-adult survival to evolve to *increase* with maternal age at early ages
319 before reversing and evolving senescence at late ages, but they emphasised that all
320 models eventually lead to accelerated rates of senescence with increasing age. This
321 prediction is consistent with the pervasive negative quadratic relationships observed
322 here, although it must be noted that Moorad and Nussey evaluated death rates on a
323 different scale than that used here in our probit models, and we cannot state with
324 confidence that a negative quadratic relationship on one scale reliably predicts the same
325 on the other. This issue can be explored explicitly in follow-up studies that can focus more
326 directly on testing evolutionary theory (see below).

327 One goal of our study was to describe the diversity of maternal ageing. In particular,
328 we were interested in the role of phylogeny. A secondary focus was to characterize how
329 human influence upon populations (e.g., laboratory vs wild populations) might affect age
330 effects. However, it was clear from our review of the relevant literature that phylogeny
331 and environment are too closely aligned to make broad conclusions regarding
332 independent effects of both factors. Consequently, we were largely forced to consider our
333 primary focus, phylogeny, but it should be understood that our suggested causal

334 inferences regarding these should be considered preliminary until enough studies that
335 disrupt the association between phylogeny and environment are published to support
336 more focused reviews. Such efforts are being made to study actuarial senescence, such as
337 in wild insect populations [63–68], and we encourage more such work to explore
338 maternal-age effects.

339 We found a reasonable number and diversity of studies (species number > 5 and
340 population replicates > 10) in laboratory invertebrates, wild birds, wild mammals, and
341 humans. Whilst maternal age in older than average individuals was clearly deleterious
342 when averaged over all groups, two taxonomic groups stood out as clearly different. Wild
343 birds were the only reasonably populated group that exhibited *positive* effects of
344 maternal age in the old. Why birds should be so different in this respect is an outstanding
345 question that requires further study, but we might speculate on possible causes.

346 Pre-adult birds are frequently cared for by individuals other than their mothers. The
347 most common source of additional care is the father [69,70]. It may be that maternal-
348 effect senescence exists in birds, but these deleterious effects are masked by sources of
349 alloparental care. Unless the ages of other care-givers are perfectly correlated with those of the
350 mothers, we can expect that offspring of old mothers will receive care from young social
351 partners and vice versa. This will obscure the effects of maternal age (an effect of a
352 reversion-to-the-mean). A total of 17 of the 20 (85%) of the surveyed wild bird species
353 are known to exhibit some form of biparental care; this approximates the prevalence of
354 80-90% across all bird species [70]. Biparental care is comparatively rare in mammals
355 [71] and invertebrates [72]. Only four of eleven surveyed wild mammal species provide
356 this form of care [71], and no surveyed invertebrate species is known to demonstrate
357 biparental care [72]. Viewed across the non-human groups studied here (invertebrates,
358 wild mammals, and wild birds), the frequency of biparental care appears to counter-

indicate the degree of maternal-effect senescence, but this pattern is only suggestive; more studies are needed from wild mammal and invertebrate species that exhibit biparental care and from bird species that do not. Interestingly, one relevant and highly-replicated laboratory study on an insect with biparental care, *Nicrophorus vespilloides*, found no effects of maternal age on pre-adult survival [19]. Furthermore, we note that even when biparental care is absent, paternal age can still have an effect on offspring outcomes through other mechanisms, such as sperm quality. For example, increased paternal age in the long-lived houbara bustard (*Chlamydotis undulata*) causes both a decline in hatching success and rate of pre-adult development [73,74]. Studies should account for variation in paternal age either by reducing or eliminating it via the experimental design (e.g., [19]) or accounting for it statistically by model fitting (e.g., [75]). Observations from human populations do not support biparental care as a primary cause for the wild bird results. Father and grandmothers can contribute meaningfully to the performance of infants and children, but humans appear to have strong signatures of maternal senescence when compared to other vertebrates, the majority of which provide only uniparental care. Finally, it must be noted that this suggested mechanism can only serve to reduce the apparent magnitude of maternal-age effects; it cannot reverse their direction.

In terms of the magnitude of effects, Lepidopterans (moths and butterflies) were clearly the most disparate group with extremely deleterious average effects in the old, an order of magnitude greater than the other groups combined. Even when compared to non-Lepidopteran invertebrates (which still exhibited stronger effects than vertebrates), these rates were seven-fold greater. Variation in the nature of maternal care might account for these differences. None of the studied invertebrate species delivered post-natal care, yet many vertebrate studies focused upon juvenile periods coincident to post-

natal care. If pre-natal maternal-effect senescence was stronger than post-natal senescence, then senescence in invertebrate studies would tend to be stronger. One way that this might happen is if increased age provides some mitigating benefit to the pre-adult. Learning, for example, is believed to cause increased fledgling rates with increased maternal age in seabirds [76,77], and it is difficult to imagine how increased experience can serve to improve pre-natal condition to the same degree. However, this suggestion does not explain why Lepidopterans age differently from other invertebrates. Pre- and post-natal maternal-effect senescence has been measured independently in very few studies, including seabirds [78] and burying beetles [19]; more such studies made over a diversity of species are necessary to assess general patterns of pre- vs post-natal maternal-effect senescence.

Another striking life history difference between studied vertebrate and invertebrate species is that the duration of reproductive lifespan of the former is longer than that of the latter, even when accounting for the vast differences in generation time. One way to quantify this is to take the square-root of the variance in standardized maternal age at birth, or $\sqrt{\sum_x \left(\frac{f(x)}{\bar{T}} - 1 \right)^2}$, where $f(x)$ is the fraction of new offspring attributed to mothers of that age, and \bar{T} is the mean of that distribution (see Materials and Methods). This provides a dimensionless comparative metric of the dispersion of maternal age (sigma). The medians of sigma for vertebrates and invertebrates are 0.347 and 0.178, respectively. Evolutionary theory anticipates that maternal senescence should evolve to be faster when sigma is small, although that prediction has not been made previously. Moorad and Nussey [31] showed that in stable age-structure populations, the strength of selection acting on an IGE produced by a mother of some age that acts on pre-adult survival is proportional to the probability $f(x)$, the same distribution that defines

generation time T (the mean) and the standard deviation about the standardized mean (sigma). It must be the case that when sigma is small, selection for IGEs declines more precipitously after T than when sigma is large. This leads to the prediction that maternal-age effects in the old are positively associated with this measure (senescence decreases as sigma grows). This might explain why Lepidopterans (median sigma = 0.069), which are usually considered to have reproductive lifespans that are so abbreviated as to make them nearly semelparous [79], senesce faster than other invertebrates (median sigma = 0.282). Further evidence for positive associations between sigma and rates of ageing come from within-group Spearman rank correlations: Lepidopterans (0.538, $n = 27$); Non-Lepidopteran Invertebrates (0.177, $n = 79$); Wild Birds (0.0325, $n = 37$); and Wild Mammals (0.112, $n = 12$). We note, however, that Lepidopterans are the only group with a correlation estimate that reaches significance ($P = 0.004$).

Finally, we note that the evolutionary theory [31] emphasized that selection for age-specific maternal IGEs for pre-adult survival follows entirely from mean vital rates. As vital rates for many species are now available (e.g. [80,81]), selection for maternal-age effects can be estimated directly over a diverse collection of populations. An obvious and tractable question that remains to be investigated is whether variation in selection explains the diversity of ageing rates within and among the groups identified here. No formal attempt has been made to reconcile patterns of selection with observations of actuarial or reproductive senescence on such a broad scale. However, this could be done in conjunction with the aforementioned analysis to evaluate which manifestations of ageing (maternal, actuarial, or reproductive) adhere closest to predictions made from evolutionary theory.

While we make no specific suggestions for why these might generate the particular patterns of diversity observed here, we believe that two other factors deserve to be

mentioned. First, the manner by which natural selection affects an evolutionary change is sensitive to the genetic architecture underlying the trait [82]. Quantitative genetic approaches can be applied to characterize genetic architecture in wild populations [83–86], but we are aware of no attempts to do this for age-specific maternal IGEs. Second, most relevant ageing studies measure cohort-level changes in the effects of maternal age. These reflect a combination of within- and among-individual changes. Evolutionary models (including [26,31]) focus on the former, which makes testing predictions using cohort-level measures risky. The most obvious source of among-individual change is selective disappearance [2,8,87,88]. Studies of maternal-effect senescence can quantify these effects, but this is rare. Among studies that do measure it, there does appear to be variation in its importance [19,51,89,90]. Evaluating the effects of selective disappearance is relatively straightforward (see cited examples), and all ageing studies should attempt to do so.

Previous studies have documented and attempted to explain among-species diversity in rates of actuarial and reproductive senescence [3,4,91]. This work extends this effort to another manifestation of ageing. Consistent with these earlier studies, we find obvious among-species variation in rates of ageing with clear evidence for underlying structure involving phylogeny. Whilst the causes for this structure are still unclear, we are encouraged that general patterns appear to be consistent with predictions from evolutionary theory, and we are hopeful that finer-scaled tests of this theory will shed light on the causes of variation in rates of maternal ageing. Future experimental and observational studies on maternal-effect senescence will improve our ability to explain this variation, especially when they focus upon understudied taxa (e.g., fish, reptiles and amphibians), wild populations of invertebrates, and species with life histories that appear unusual for their taxonomic groupings (such as mammals or insects that exhibit

paternal or cooperative care). Taxonomic gaps amongst wild populations likely reflect a general preference amongst ecologists to invest in long-term studies of birds and mammals rather than any biological feature of these species that might make them more amenable to the study of ageing. Finally, it would be interesting to assess maternal ageing in species that lack evidence for actuarial senescence [see 4] (these species did not appear in our search for data amenable to our analyses), particularly as evolutionary theory predicts a link between adult ageing rates and the evolution of maternal senescence [31].

Authors' Contributions

EIC and JM conceived the ideas, designed the methodology, analysed the data and wrote the manuscript. EIC collected the data. Both gave final approval for publication.

Acknowledgements

This research was supported by East of Scotland Bioscience Doctoral Training Partnership (EASTBIO) and the Biotechnology and Biological Sciences Research Council (BBSRC) (544EIC BB/J01446X/1). We are grateful to Katie McGhee, Jean-Michel Gaillard, Marco Festa-Bianchet, and one anonymous reviewer for valuable comments on the manuscript. Also to Ally Phillimore, Jarrod Hadfield, Joshua Moatt, Liam Brierly, Alex Scheuerlein, Joel Pick, and members of the Moorad and Smiseth lab groups for helpful advice and comments. The authors declare no conflict of interest.

References

1. Descamps S, Boutin S, Berteaux D, Gaillard J-M. 2008 Age-specific variation in survival, reproductive success and offspring quality in red squirrels: evidence of senescence. *Oikos* **117**, 1406–1416. (doi:10.1111/j.2008.0030-1299.16545.x)

- 483 2. Bouwhuis S, Sheldon BC, Verhulst S, Charmantier A. 2009 Great tits growing old:
484 selective disappearance and the partitioning of senescence to stages within the
485 breeding cycle. *Proc. R. Soc. London B Biol. Sci.* **276**, 2769–77.
486 (doi:10.1098/rspb.2009.0457)
- 487 3. Jones OR *et al.* 2008 Senescence rates are determined by ranking on the fast-slow
488 life-history continuum. *Ecol. Lett.* **11**, 664–673. (doi:10.1111/j.1461-
489 0248.2008.01187.x)
- 490 4. Jones OR *et al.* 2014 Diversity of ageing across the tree of life. *Nature* **505**, 169.
491 (doi:10.1038/nature12789)
- 492 5. Bonduriansky R, Brassil CE. 2002 Rapid and costly ageing in wild male flies.
493 *Nature* **420**, 377. (doi:10.1038/420377a)
- 494 6. Waugh SM, Barbraud C, Adams L, Freeman AND, Wilson K-J, Wood G, Landers TJ,
495 Baker GB. 2015 Modeling the demography and population dynamics of a
496 subtropical seabird, and the influence of environmental factors. *Condor* **117**, 147–
497 164. (doi:doi:10.1650/CONDOR-14-141.1)
- 498 7. Nussey DH, Coulson T, Festa-Bianchet M, Gaillard J-M. 2008 Measuring
499 senescence in wild animal populations: Towards a longitudinal approach. *Funct.*
500 *Ecol.* **22**, 393–406. (doi:10.1111/j.1365-2435.2008.01408.x)
- 501 8. Nussey DH, Coulson T, Delorme D, Clutton-Brock TH, Pemberton JM, Festa-
502 Bianchet M, Gaillard J-M. 2011 Patterns of body mass senescence and selective
503 disappearance differ among three species of free-living ungulates. *Ecology* **92**,
504 1936–1947. (doi:10.1890/11-0308.1)
- 505 9. Gaillard J-M, Allaine D, Pontier D, Yoccoz NG, Promislow DEL. 1994 Senescence in
506 natural populations of mammals: A reanalysis. *Evolution (N. Y.)* **48**, 509–516.
507 (doi:10.2307/2410110)

- 508 10. Lemaître J-F, Gaillard J-M. 2017 Reproductive senescence: new perspectives in
509 the wild. *Biol. Rev.* **92**, 2182–2199. (doi:10.1111/brv.12328)
- 510 11. Rose MR. 1984 Laboratory evolution of postponed senescence in *Drosophila*
511 *melanogaster*. *Evolution (N. Y.)*. **38**, 1004–1010. (doi:10.2307/2408434)
- 512 12. Kenyon C, Chang J, Gensch E, Rudner A, Tabtiang R. 1993 A *C. elegans* mutant that
513 lives twice as long as wild type. *Nature* **366**, 461–464. (doi:10.1038/366461a0)
- 514 13. Bonduriansky R, Maklakov A, Zajitschek F, Brooks R. 2008 Sexual selection, sexual
515 conflict and the evolution of ageing and life span. *Funct. Ecol.* **22**, 443–453.
516 (doi:10.1111/j.1365-2435.2008.01417.x)
- 517 14. Galliot B. 2012 Hydra, a fruitful model system for 270 years. *Int. J. Dev. Biol.* **56**,
518 411–423. (doi:10.1387/ijdb.120086bg)
- 519 15. Bitton PP, Dawson RD. 2017 Age-related prenatal maternal effects and postnatal
520 breeding experience have different influences on nestling development in an
521 altricial passerine. *J. Avian Biol.* **48**, 660–668. (doi:10.1111/jav.01202)
- 522 16. Clark J, Garbutt JS, McNally L, Little TJ. 2017 Disease spread in age structured
523 populations with maternal age effects. *Ecol. Lett.* **20**, 445–451.
524 (doi:10.1111/ele.12745)
- 525 17. Lippens C, Faivre B, Lechenault C, Sorci G. 2017 Aging parasites produce offspring
526 with poor fitness prospects. *Biol. Lett.* **13**, 20160888.
527 (doi:10.1098/rsbl.2016.0888)
- 528 18. Bouwhuis S, Vedder O, Becker PH. 2015 Sex-specific pathways of parental age
529 effects on offspring lifetime reproductive success in a long-lived seabird.
530 *Evolution (N. Y.)*. **69**, 1760–1771. (doi:10.1111/evo.12692)
- 531 19. Ivimey-Cook ER, Moorad J. 2018 Disentangling pre- and post-natal maternal age
532 effects on offspring performance in an insect with elaborate maternal care. *Am.*

- 533 *Nat.* **192**, 564–576.
- 534 20. Bloch Qazi MC, Miller PB, Poeschel PM, Phan MH, Thayer JL, Medrano CL. 2017
- 535 Transgenerational effects of maternal and grandmaternal age on offspring
- 536 viability and performance in *Drosophila melanogaster*. *J. Insect Physiol.* **100**, 43–
- 537 52. (doi:10.1016/j.jinsphys.2017.05.007)
- 538 21. Harman D. 1956 Aging: A theory based on free radical and radiation chemistry. *J.*
- 539 *Gerontol.* **11**, 298–300. (doi:10.1093/geronj/11.3.298)
- 540 22. Eichhorn GL, Shin YA, Clark P, Rifkind J, Pitha J, Tarien E, Rao G, Karlik SJ, Crapper
- 541 DR. 1979 Essential and deleterious effects in the interaction of metal ions with
- 542 nucleic acids. *Trace Met. Heal. Dis.* , 123–134.
- 543 23. Medvedev ZA. 1986 Age-related changes of transcription and RNA processing. In
- 544 *Drugs and Aging* (ed. D. Platt), pp. 1–19. Berlin, Heidelberg: Springer Berlin
- 545 Heidelberg. (doi:10.1007/978-3-642-70788-9_1)
- 546 24. Orgel LE. 1970 The maintenance of the accuracy of protein synthesis and its
- 547 relevance to ageing: a correction. *Proc. Natl. Acad. Sci. U. S. A.* **67**, 1476.
- 548 (doi:10.1073/pnas.67.3.1476)
- 549 25. Medvedev ZA. 1990 An attempt at a rational classification of theories of ageing.
- 550 *Biol. Rev.* **65**, 375–398. (doi:10.1111/j.1469-185X.1990.tb01428.x)
- 551 26. Hamilton WD. 1966 The moulding of senescence by natural selection. *J. Theor.*
- 552 *Biol.* **12**, 12–45. (doi:10.1016/0022-5193(66)90184-6)
- 553 27. Medawar PB. 1952 *An Unsolved Problem Of Biology*. London: Lewis.
- 554 28. Williams GC. 1957 Pleiotropy, natural selection, and the evolution of senescence.
- 555 *Evolution (N. Y.)*. **11**, 398–411. (doi:10.2307/2406060)
- 556 29. Hughes KA, Charlesworth B. 1994 A genetic analysis of senescence in *Drosophila*.
- 557 *Nature* **367**, 64–66. (doi:10.1038/367064a0)

- 558 30. Charlesworth B. 2001 Patterns of age-specific means and genetic variances of
559 mortality rates predicted by the mutation-accumulation theory of ageing. *J. Theor.*
560 *Biol.* **210**, 47–65. (doi:10.1006/jtbi.2001.2296)
- 561 31. Moorad JA, Nussey DH. 2016 Evolution of maternal effect senescence. *Proc. Natl.*
562 *Acad. Sci. U. S. A.* **113**, 362–367. (doi:10.1073/pnas.1520494113)
- 563 32. Moher D, Liberati A, Tetzlaff J, Altman DG, Grp P. 2009 Preferred reporting items
564 for systematic reviews and meta-analyses: the PRISMA statement. *Phys. Ther.* **89**,
565 873–880. (doi:10.1371/journal.pmed.1000097)
- 566 33. Charlesworth B. 1994 *Evolution in Age-Structured Populations*. Cambridge:
567 Cambridge University Press. See
568 [https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1685027/pdf/ajhg00363-](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1685027/pdf/ajhg00363-0169.pdf)
569 [0169.pdf](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1685027/pdf/ajhg00363-0169.pdf).
- 570 34. Packer C, Tatar M, Collins A. 1998 Reproductive cessation in female mammals.
571 *Nature* **392**, 807–11. (doi:10.1038/33910)
- 572 35. Fowler K, Partridge L. 1989 A cost of mating in female fruitflies. *Nature* **338**, 760–
573 761. (doi:10.1038/338760a0)
- 574 36. Ashburner M, Thompson JN (1978). The laboratory culture of *Drosophila*. In: The
575 genetics and biology of *Drosophila*. (Ashburner M, Wright TRF (eds.)). Academic
576 Press, volume 2A: pp. 1–81.
- 577 37. Federhen S. 2011 The NCBI Taxonomy. *Nucleic Acids Res.* **40**, D136–D143.
578 (doi:10.1093/nar/gkr1178)
- 579 38. Letunic I. 2015 phyloT: phylogenetic Tree Generator. Phylot.biobyte.de. *Nucl.*
580 *Acids Res.* **10**, 725, (doi:10.1093/nar/gkw290)
- 581 39. Wickham H. 2009 *ggplot2 Elegant Graphics for Data Analysis*. (doi:10.1007/978-
582 0-387-98141-3)

- 583 40. Yu G, Smith DK, Zhu H, Guan Y, Lam TTY. 2017 ggtree: an r package for
584 visualization and annotation of phylogenetic trees with their covariates and other
585 associated data. *Methods Ecol. Evol.* **8**, 28–36. (doi:10.1111/2041-210X.12628)
- 586 41. Moorad JA, Promislow DEL. 2011 Evolutionary demography and quantitative
587 genetics: Age-specific survival as a threshold trait. *Proc. R. Soc. B Biol. Sci.* **278**,
588 144–151. (doi:10.1098/rspb.2010.0992)
- 589 42. Robertson A, Lerner IM. 1949 The heritability of all-or-none traits; viability of
590 poultry. *Genetics* **34**, 395.
- 591 43. Hurvich CM, Tsai CL. 1989 Regression and time series model selection in small
592 samples. *Biometrika* **76**, 297–307. (doi:10.1093/biomet/76.2.297)
- 593 44. Kushary D, Davison AC, Hinkley D V. 2000 Bootstrap methods and their
594 application. *Technometrics* **42**, 216. (doi:10.2307/1271471)
- 595 45. Canty A, Ripley B. 2017 boot: Bootstrap R (S-Plus) functions. *R Packag. version*
596 *1.3-20* (doi:10.1007/978-1-4614-7618-4)
- 597 46. R Core Team. 2016 R: A language and environment for statistical computing. *R*
598 *Found. Stat. Comput. Vienna Austria.* **0**, {ISBN} 3-900051-07-0.
599 (doi:10.1038/sj.hdy.6800737)
- 600 47. Gilbert JJ. 2010 Effect of food concentration on the production and viability of
601 resting eggs of the rotifer *Brachionus*: Implications for the timing of sexual
602 reproduction. *Freshw. Biol.* **55**, 2437–2446. (doi:10.1111/j.1365-
603 2427.2010.02455.x)
- 604 48. Alonso-Alvarez C, Perez-Rodriguez L, Garcia JT, Vinuela J, Mateo R. 2010 Age and
605 breeding effort as sources of individual variability in oxidative stress markers in a
606 bird species. *Physiol. Biochem. Zool.* **83**, 110–118. (doi:10.1086/605395)
- 607 49. Mourão JL, Barbosa AC, Outor-Monteiro D, Pinheiro VM. 2010 Age affects the

608 laying performance and egg hatchability of red-legged partridges (*Alectoris rufa*)
609 in captivity. *Poult. Sci.* **89**, 2494–2498. (doi:10.3382/ps.2009-00377)

610 50. Annett RW, Carson AF, Dawson LER, Irwin D, Kilpatrick DJ. 2011 Effects of breed
611 and age on the performance of crossbred hill ewes sourced from Scottish
612 Blackface dams. *Animal* **5**, 356–366. (doi:10.1017/S1751731110002090)

613 51. Hayward AD, Wilson AJ, Pilkington JG, Clutton-Brock TH, Pemberton JM, Kruuk
614 LEB. 2013 Reproductive senescence in female Soay sheep: Variation across traits
615 and contributions of individual ageing and selective disappearance. *Funct. Ecol.*
616 **27**, 184–195. (doi:10.1111/1365-2435.12029)

617 52. Skibił AL, Dobson FS, Murie JO. 2009 Maternal influences on reproduction in two
618 populations of Columbian ground squirrels. *Ecol. Monogr.* **79**, 325–341.
619 (doi:10.1890/08-0718.1)

620 53. Neuhaus P, Broussard DR, Murie JO, Dobson FS. 2004 Age of primiparity and
621 implications of early reproduction on life history in female Columbian ground
622 squirrels. *J. Anim. Ecol.* **73**, 36–43. (doi:10.1111/j.1365-2656.2004.00793.x)

623 54. Lemaître J-F, Gaillard J-M, Lackey LB, Clauss M, Müller DWH. 2013 Comparing
624 free-ranging and captive populations reveals intra-specific variation in aging
625 rates in large herbivores. *Exp. Gerontol.* **48**, 162–167.
626 (doi:10.1016/j.exger.2012.12.004)

627 55. Promislow DEL. 1991 Senescence in natural populations of mammals: a
628 comparative study. *Evolution (N. Y.)*. **4**, 1869–1887. (doi:10.2307/2409837)

629 56. Tidière M, Gaillard J-M, Berger V, Müller DWH, Bingaman Lackey L, Gimenez O,
630 Clauss M, Lemaître J-F. 2016 Comparative analyses of longevity and senescence
631 reveal variable survival benefits of living in zoos across mammals. *Sci. Rep.* **6**,
632 36361. (doi:10.1038/srep36361)

- 633 57. Griffing B. 1968 Selection in reference to biological groups. 3. Generalized results
634 of individual and group selection in terms of parent-offspring covariances. *Aust. J.*
635 *Biol. Sci.* **21**, 1171–1178. (doi:10.1071/BI9681171)
- 636 58. Griffing B. 1967 Selection in reference to biological groups. I. Individual and
637 group selection applied to populations of unordered groups. *Aust. J. Biol. Sci.* **20**,
638 127–139. (doi:10.1071/bi9670127)
- 639 59. Moore AJ, Brodie III ED, Wolf JB. 1997 Interacting phenotypes and the
640 evolutionary process: I. Direct and indirect genetic effects of social interactions.
641 *Evolution (N. Y.)*. **51**, 1352–1362. (doi:10.2307/2411187)
- 642 60. Wolf JB, Brodie III ED, Moore AJ. 1999 Interacting phenotypes and the
643 evolutionary process. II. Selection resulting from social interactions. *Am. Nat.* **153**,
644 254–266. (doi:10.1086/303168)
- 645 61. McGlothlin JW, Moore AJ, Wolf JB, Brodie ED. 2010 Interacting phenotypes and
646 the evolutionary process. III. Social evolution. *Evolution (N. Y.)*. **64**, 2558–2574.
647 (doi:10.1111/j.1558-5646.2010.01012.x)
- 648 62. Kirkwood TBL. 1977 Evolution of ageing. *Nature* **270**, 301–304.
- 649 63. Kawasaki N, Brassil CE, Brooks RC, Bonduriansky R. 2008 Environmental effects
650 on the expression of life span and aging: an extreme contrast between wild and
651 captive cohorts of *Telostylinus angusticollis* (Diptera: Neriidae). *Am. Nat.* **172**,
652 346–357. (doi:10.1086/589519)
- 653 64. Carroll J, Sherratt TN. 2017 Actuarial senescence in laboratory and field
654 populations of Lepidoptera. *Ecol. Entomol.* **42**, 675–679.
655 (doi:10.1111/een.12417)
- 656 65. Sherratt TN, Laird RA, Hassall C, Lowe CD, Harvey IF, Watts PC, Cordero-Rivera A,
657 Thompson DJ. 2010 Empirical evidence of senescence in adult damselflies

- (Odonata: Zygoptera). *J. Anim. Ecol.* **79**, 1034–1044. (doi:10.1111/j.1365-2656.2010.01719.x)
66. Mautz BS, Rode NO, Bonduriansky R, Rundle HD. 2019 Comparing ageing and the effects of diet supplementation in wild vs. captive antler flies, *Protopiophila litigata*. *J. Anim. Ecol.* **88**, 1913–1924. (doi:10.1111/1365-2656.13079)
67. Rodríguez-Muñoz R, Boonekamp JJ, Liu XP, Skicko I, Haugland Pedersen S, Fisher DN, Hopwood P, Tregenza T. 2019 Comparing individual and population measures of senescence across 10 years in a wild insect population. *Evolution (N. Y.)*. **73**, 293–302. (doi:10.1111/evo.13674)
68. Zajitschek F, Zajitschek S, Bonduriansky R. 2020 Senescence in wild insects: Key questions and challenges. *Funct. Ecol.* **34**, 26–37. (doi:10.1111/1365-2435.13399)
69. Lack D. 1968 *Ecological adaptations for breeding in birds*. London, UK: Methuen & Co. See <http://agris.fao.org/agris-search/search.do?recordID=US201300591006>.
70. Cockburn A. 2006 Prevalence of different modes of parental care in birds. *Proc. R. Soc. B Biol. Sci.* **273**, 1375–1383. (doi:10.1098/rspb.2005.3458)
71. Stockley P, Hobson L. 2016 Paternal care and litter size coevolution in mammals. *Proc. R. Soc. B Biol. Sci.* **283**, 20160140. (doi:10.1098/rspb.2016.0140)
72. Suzuki S. 2014 Biparental care in insects: Paternal care, life history, and the function of the nest. *J. Insect Sci.* **13**, 1–16. (doi:10.1673/031.013.13101)
73. Preston BT, Saint Jalme M, Hingrat Y, Lacroix F, Sorci G. 2015 The sperm of aging male bustards retards their offspring's development. *Nat. Commun.* **6**, 1–9. (doi:10.1038/ncomms7146)
74. Evans JP, Wilson AJ, Pilastro A, Garcia-Gonzalez F. 2019 Ejaculate-mediated paternal effects: Evidence, mechanisms and evolutionary implications.

683 *Reproduction*. **153**, R109–R126. (doi:10.1530/REP-18-0524)

684 75. Fay R, Barbraud C, Delord K, Weimerskirch H. 2016 Paternal but not maternal age
685 influences early-life performance of offspring in a long-lived seabird. *Proc. R. Soc.
686 B Biol. Sci.* **283**, 20152318. (doi:10.1098/rspb.2015.2318)

687 76. Froy H, Phillips RA, Wood AG, Nussey DH, Lewis S. 2013 Age-related variation in
688 reproductive traits in the wandering albatross: evidence for terminal
689 improvement following senescence. *Ecol. Lett.* **16**, 642–649.
690 (doi:10.1111/ele.12092)

691 77. Part T. 1995 Does breeding experience explain increased reproductive success
692 with age? An experiment. *Proc. R. Soc. B Biol. Sci.* **260**, 113–117.
693 (doi:10.1098/rspb.1995.0067)

694 78. Beamonte-Barrientos R, Velando A, Drummond H, Torres R. 2010 Senescence of
695 maternal effects: aging influences egg quality and rearing capacities of a long-
696 lived bird". *Am. Nat.* **175**, 469–470. (doi:10.1086/650726)

697 79. Fritz RS, Stamp NE, Halverson TG. 1982 Iteroparity and semelparity in insects.
698 *Am. Nat.* **120**, 264–268. (doi:10.2307/2678832)

699 80. Max-Planck Institute for Demographic Research. In press. DATLife Database. See
700 www.datlife.org (accessed on 3 March 2019).

701 81. Salguero-Gomez R *et al.* 2016 COMADRE: a global database of animal
702 demography. *J. Anim. Ecol.* **85**, 371–384. (doi:10.1111/027821)

703 82. Lande R, Arnold SJ. 1983 The measurement of selection on correlated characters.
704 *Evolution (N. Y.)*. **37**, 1210. (doi:10.2307/2408842)

705 83. Charmantier A, Perrins C, McCleery RH, Sheldon BC. 2006 Quantitative genetics of
706 age at reproduction in wild swans: Support for antagonistic pleiotropy models of
707 senescence. *Proc. Natl. Acad. Sci.* **103**, 6587–6592.

- (doi:10.1073/pnas.0511123103)
84. McCleery RH, Perrins CM, Sheldon BC, Charmantier A. 2008 Age-specific reproduction in a long-lived species- the combined effects of senescence and individual quality. *Proc. R. Soc. B* **275**, 963–970. (doi:10.1098/rspb.2007.1418)
85. Nussey DH, Wilson AJ, Morris A, Pemberton J, Clutton-Brock T, Kruuk LEBB. 2008 Testing for genetic trade-offs between early- and late-life reproduction in a wild red deer population. *Proc. R. Soc. B Biol. Sci.* **275**, 745–750. (doi:10.1098/rspb.2007.0986)
86. Wilson AJ, Pemberton JM, Pilkington JG, Clutton-Brock TH, Coltman DW, Kruuk LEB. 2007 Quantitative genetics of growth and cryptic evolution of body size in an island population. *Evol. Ecol.* **21**, 337–356. (doi:10.1007/s10682-006-9106-z)
87. Vaupel JW, Manton KG, Stallard E. 1979 The impact of heterogeneity in individual frailty on the dynamics of mortality. *Demography* **16**, 439–454. (doi:10.2307/2061224)
88. Vaupel JW, Yashin AI. 1985 Heterogeneity's ruses: Some surprising effects of selection on population dynamics. *Am. Stat.* **39**, 176–185. (doi:10.1080/00031305.1985.10479424)
89. Zhang H, Vedder O, Becker PH, Bouwhuis S. 2015 Age-dependent trait variation: The relative contribution of within-individual change, selective appearance and disappearance in a long-lived seabird. *J. Anim. Ecol.* **84**, 797–807. (doi:10.1111/1365-2656.12321)
90. van de Pol M, Verhulst S. 2006 Age-dependent traits: a new statistical model to separate within- and between-individual effects. *Am. Nat.* **167**, 766–773. (doi:10.1086/503331)
91. Holmes DJ, Austad SN. 1995 The evolution of avian senescence patterns:

Implications for understanding primary aging processes. *Integr. Comp. Biol.* **35**,
307–317. (doi:10.1093/icb/35.4.307)

Figure Legends

Fig. 1 Maternal-age effects in old individuals arranged by phylogenetic relationship. Effect sizes and SEs (on the probit scale) were averaged within species. Three replicates were removed prior to averaging as their SEs made visualisation impossible (*Centropages typicus*, -78.53 ± 8991.38 ; *Psuedaletia sequax*, -159.74 ± 6298.32 ; and *Ovis aries*, 14.23 ± 2474.36). Two species indicated by * were not included here because their estimates would not fit (see Table S3b for these). Colours/line-type indicate environment: wild (blue/two-dashes), laboratory (red/dot-dash), semi-captive (purple/long-dash), agricultural (black/solid), and humans (orange/dotted). Error bars around the estimate represent 95% confidence intervals.